Estuaries Vol. 3, No. 4, p. 329–333 December 1980

General Index

A

Ablabesmyia spp., 50
Acanthohaustorius
  bousfieldi new species, 75–79, 82
  key to species, 75–76
  millsi, 79, 82–83
  shoemakeri, 79
  similis new species, 75, 79–83
  spinosus, 79
  sp., 79
Accipiter
  gentilis, 165
  nisus, 159
Acipenser oxyrninchus, 38
Acrochaetium, 204
Adams Cove, New Hampshire, 28–37
Agardhiella tenera, 204
Agrostis stolonifera, 174
  tenuis, 174
algae, effect of atrazine on, 271–277
Algal
  epiphyte productivity, 315–317
  mat productivity, 122–131
Alisma spp., 173
Almyracuma proximoculi, 42, 50
Alosa
  aestivalis, 52, 65–68
  pseudoharengus, 52
  sapidissima, 52, 65–68
Alpheus heterochaelis, 293
American oyster, 186, 189
amphipods, new species, 75–83
Anas
  platyrhynchos, 163
  rubripes, 165
Anchoa
  hepsetus, 92
  mitchilli, 278, 280
Anderson, C. M., and M. Treshow, 168–176
Anderson, F. E., 28–37
Apalachee Bay, Florida, 177–183
Apeltes quadracus, 281
Ardea herodias, 164
Arellano, E., and W. S. Brown, 248–257
arrowleaf, 173
arsenic, 155, 157, 164
Artemia salina, 321–323
Ascophyllum, 200, 205
Asiatic clam, 296
Asterionella glacialis, 208, 213, 215
Atrazine effects
  on algae, 271–277
  on *Uca pugnax*, 217–223
Aythya valisineria, 162

B

bacteria, density on decomposing *Spartina*, 61–64
Bairdiella chrysoura, 278, 280
bald eagle, 156
Barataria Bay, Louisiana, 112–121
Barker, H. R., Jr., *see* Dauer, D. M.
Batis maritima, 122–129
behavioral responses, to naphthalene by blue crab, 224–229
benthos
  in *Myriophyllum spicatum* beds, 38–54
  of Sacramento River, 296–307
Bidleman, T. F., *see* Harder, H. W.
bird
  barn owl, 164
  black duck, 165
  Canada geese, 163
  canvasback, 162
  cowbird, 164
  European kestrel, 165
  great blue heron, 164
  goshawk, 165
  lesser flamingo, 164
  mallard, 163, 165
  pheasant, 165
  red-tailed hawk, 165
  ruffed grouse, 162
  tawny owl, 164
  white pelican, 164
Boccardia ligerica, 296, 301–304
Bonasa umbellus, 162
Bostrychia radicans, 203
Bozeman, E. L., Jr., and J. M. Dean, 89–97
Branta canadensis, 163
breeding beaches, of horseshoe crab, 177–183
Brevoortia tyrannus, 89, 93–95, 189, 282
brown pelican, 162
Brown, W. S., and E. Arellano, 248–257
Buresh, R. J., R. D. DeLanne and W. H. Patrick, Jr., 111–121
Burnett, L. E., *see* Middaugh, D. P.
Buteo jamaicensis, 165

C

cadmium, 155, 157, 164
Callinectes sapidis, 146, 224–229, 282, 286, 289–290, 292, 293
  detection threshold to naphthalene, 224–229
  kepone in, 192–199
Callithamnion byssoides, 203
Caloglossa leprieurii, 203
Cammen, L. M., 55–60
Cape Fear River, North Carolina, 98–105
carbon
  dissolved organic released by phytoplankton, 230–233
  fixation, 315–317
Carcharhinus milberti, 278–286
Carcinoscorpus rotundicauda, 182
Cardium edule, 84
Cargo, D. G., and G. E. Rabenold, 20–27
cattail, broad-leaved, 173
Centropristis striata, 92, 280
Ceramium, 200, 203, 205
  areschougii, 204
  deslongchampii, 204
  diaphanum, 204
  rubriforme, 204
  rubrum, 204
  strictum, 204
Cerataulina pelagica, 208, 212, 215
Ceratium furca, 213, 215
Cerithidea californica, 127
Chaetoceros
  spp., 231–232
  sociale, 215
Chaetognaths, 106–110
Chasmodes bosquianus, 280
Chesapeake Bay, 84–87, 200–206, 207–216, 242–247, 278–288, 289–294
Chiridotea almyra, 42
Chironomid population of Hudson River, 38–54
Chironomus decorus, 44–47, 49
  spp., 50
Chlordanes, 155, 159
Chlorella pyrenoidosa, 275
Chlorinated sewage effluent, detoxification, 184–191
Chlorine produced oxidants, response of fish, 132–141
Chondria sp., 204
Christensen, E. C., *see* Harder, H. W.
Chromium, 155, 157, 162
Chroomonas caroliniana, 213
Chrysaora quinquecirrha, Asexual reproduction, 20–27
Ciliates, density on decomposing *Spartina*, 61–64
Circulation and mixing, in Potomac River Estuary, 11–19
Cladophora, 200, 205
Clam
  Asiatic, 150–151
  soft shell, 84
Clayton, G. R., *see* Murawski, S. A.
Climacodium biconcavum, 215
Cochlodinium helicoides, 187, 189
Coelotanypus scapularis, 50
Cole, C. F., *see* Murawski, S. A.
Collection efficiency, seine and rotenone, 98–105
Congeria leucophaeta, 49
Connecticut River, 65–68
Copper, 157, 163
Corbicula
  fluminea, 150–151
  manilensis, 296, 301–305
Cordylophora lacustris, 49
Corophium
  lacustre, 50
  spinicorne, 296, 301–303
  stimpsoni, 296, 301–305
Cory, R. L., and P. V. Dresler, 150–151
Coscinodiscus marginatus, 208, 215
Couch, J. A., *see* Middaugh, D. P.
Crab
  blue, 224–229, 282, 286, 289
    Kepone in, 192–199

fiddler, 217–223, 272–274
hermit, 290
mud, 290
shore, 228
spider, 290
Crangon
franciscorum, 301
septemspinosa, 1–10, 289, 290, 292
Crassostrea virginica, 146, 184, 186, 189
Cricotopus sylvestris, 41, 44–53
sp., 50
Cryptochironomus nr. *fulvus*, 50
Culture system for mysid shrimps, 321–323
Cyathura polita, 42, 50
Cyclotella
caspia, 214
glomerata, 208, 212
meneghiniana, 276
Cymatosira belgica, 271, 274–275
Cynoscion regalis, 282
Cyprinodon variegatus, 228

D

Daphnia magna, 185
Dauer, D. M., R. M. Ewing, G. H. Tourtellotte, and H. R. Barker, Jr., 148–149
Davis, D. E., and F. G. Plumley, 271–277
see Plumley, F. G.
Davis, R. W., and M. P. Weinstein, 98–105
DDD, 155, 159
DDE, 155, 159
DDT, 155–156, 159
Dean, J. M., and E. L. Bozeman, Jr., 89–97
Decapod crustacea, in eelgrass meadows, 289–295
dechlorination of sewage effluent, 184–191
Decomposition rate, of *Spartina alterniflora*, 61–64
DeLaune, R. D., *see* Buresh, R. J.
D'Elia, C. F., Book review, 152
Denticula subtilis, 125
Deposit feeders, measuring ingestion rate, 55–60
Desiccation
effects on algal productivity, 315–317
tolerance of *Ilyanassa obsoleta*, 318–320
Desmotrichum undulatum, 200, 205
Detection threshold
of blue crab for naphthalene, 224–229
Diatom
density on decomposing *Spartina*, 61–64
growth rate, 69–72
Dicrotendipes modestus, 44, 45, 47–48, 50–51
nervosus, 50
Dieldrin, 155–156, 159
Distichlis spicata, 266
Dormermuth, R. B., and R. J. Reed, 65–68
Dresler, P. V., and R. L. Cory, 150–151
Drewry, G. E., book review, 73–74

E

Ectocarpus confervoides, 200
spp., 204–205
Eelgrass meadows, in Chesapeake Bay, 278–288, 289–295
Eggs, blue crab, 192–199
Eichornia crassipes, 173
Enallagma durum, 49, 52
Enteromorpha, 174
clathrata var. *crinita*, 125
compressa, 200, 205
linza, 200, 205
Entocladia viridis, 200, 205
Epifauna, on *Myriophyllum spicatum*, 38–54
Erosion in tidal marshes, 258–270
European sparrowhawk, 159
Eurytium spp., 220
Everest, J. W., *see* Plumley, F. G.
Ewing, R. M., *see* Dauer, D. M.

F

Falco tinnunculus, 165
Fish
alewife, 39, 52
American shad, 52, 65–68
Atlantic menhaden, 93, 189
Atlantic croaker, 92
Atlantic silverside, 103, 186–187, 280
Atlantic sturgeon, 38
bay anchovy, 278
bighead searobin, 92
biomass, 91–95
blackcheek tongue fish, 92
black sea bass, 92, 280
blueback herring, 52, 65–68
bluegill, 38, 51, 140
brook trout, 142
chain pipefish, 92
chum salmon, 313
coho salmon, 140
fathead minnow, 133, 142
four-spined stickleback, 281
golden shiner, 185
grunion, 177
larval and juvenile abundance, 89–97
longfin smelt, 308–314
mosquitofish, 140
mullet, 99–104
mummichog, 146
northern pipefish, 278–286
pink salmon, 313
pinfish, 93, 278–286, 292
pumkinseed, 65–68
rainbow smelt, 308–314
rainbow trout, 133, 140
sandbar shark, 278–286
silver perch, 278, 280–286
silverside, 99–104
speckled worm eel, 93
spot, 92, 99–104, 189, 278–286
response to chlorine produced oxidants, 132–141
spottail shiner, 39
striped anchovy, 92
striped mullet, 93
striped bass, 51, 133
summer flounder, 280
tautog, 280
white mullet, 93
white perch, 38, 139–140
Fish, food of
alewife, 39, 52
American shad, 52, 65–68
Atlantic sturgeon, 38
blueback herring, 52, 65–68
bluegill, 38, 51
pumpkinseed, 65–68
spottail shiner, 39
white perch, 38
Flagellates, density on decomposing *Spartina*, 61–64
Frame, A. B., 75–83
Fresh water discharge, into Great Bay Estuary, 248–257
Freezing tolerance of *Ilyanassa obsoleta*, 318–320
Fundulus heteroclitus, 146, 228, 292
Fungi, density on decomposing *Spartina*, 61–64

G

Gambusia affinis, 140
Gammarus daiberi, 49–50, 52
Garofalo, D., 258–270
Geukensia (=*Modiolus*) *demissa*, 84–87, 218
Gracilaria foliifera, 204
verrucosa, 204
Grass shrimp, 198
Great Bay Estuary, New Hampshire, 248–257
Grinnelia americana, 200, 203
Growth rate of diatom, 69–72
Gulf of Mexico, 84–87
Gymnodinium danicans, 208, 212
Gyrodinium estuariale, 208, 212
Gyrosigma obliquum, 125

H

Haliaeetus leucocephalus, 156
Hamilton, A. D., and R. E. Wilson, 11–19
Harder, H. W., E. C. Christensen, J. R. Matthews, and T. F. Bideman, 142–147
Hartwig, E. O., *see* Taft, J. L.
Heck, K. L., Jr., and R. J. Orth, 278–288, 289–295
Heptachlor epoxide, 155–159
Hippolyte pleuracantha, 293
Holyoke Dam, Massachusetts, 65–68
Horseshoe crab, breeding behavior and movements, 177–183
Hudson River Estuary, 38–54
Hydra sp., 49

Hydrography, of a tidal creek, 236–241
Hypaniola florida, 42
Hypsoblennius hentzi, 280

I

Ilyanassa obsoleta, 318–320
Ingestion rate of deposit feeders, 55–60
Inundation frequency, effect on algal productivity, 315–317
Ischadium recurvum, 85–87

J

Jaumea carnosa, 122–129
Jones Creek, South Carolina, 236–241
Jones, R. C., 315–317
Juncus roemerianus, 240

K

Kepone in blue crab eggs and muscle, 192–199
Kjerfve, B., and F. B. Schwing, 236–241
Knight, A. W., *see* Siegfried, C. A.
Kopache, M. E., *see* Siegfried, C. A.

L

Lafayette River, Virginia, 148–149
Lagodon rhomboides, 89, 93–95, 278–286, 292
Lamont, T. G., *see* Wiemeyer, S. N.
Lead, 155, 157, 165
Leggett, A. T., Jr., and M. H. Roberts, Jr., 192–199
Leiostomus xanthurus, 89, 92–95, 99–104, 189, 190, 278–286
  response to chlorine produced oxidants, 132–141
Lepomis
  gibbosus, 65–68
  macrochirus, 39, 51, 140
Leptochierus plumosus, 50
Leptocylindrus danicus, 213, 215
Leuresthes tenuis, 177
Libinia dubia, 290
Life cycle, *Crangon septemspinosa*, 1–10
Limnodrilus
  hoffmeisteri, 51
  profundicola, 51
Limulus polyphemus, 177–183
Littorina irrorata, 218
Locke, L. N., *see* Wiemeyer, S. N.
Loftus, R., *see* Taft, J. L.
Lomentaria baileyana, 203
Long Island Sound, 75–83

M

Macoma balthica, 296, 303
Macroalgae of Chesapeake Bay, 200–206
Marshall, H. G., 207–216
Matthews, J. R., *see* Harder, H. W.
McCausland, E., and D. J. Murphy, 318–320
McEnerney, *see* Plumley, F. G.
Melosira nummuloides, growth response to UV-B radiation 69–72
Menidia
  menidia, 103, 184, 186–187, 189, 280
  spp., 99
Menzie, C. A., 38–54
Mercury, 155, 157, 164–165
Microcoleus lyngbyaceus, 125
Micropogon undulatus, 89, 92–95
Micropogonias undulatus, 282
Middaugh, D. P., L. E. Burnett, and J. A. Couch, 132–141
Migration of tidal stream channels, 258–270
Model, segmented tidal mixing, 248–257
Modlin, R. F., 1–10
Molothrus ater, 164
Monanthochloe littoralis, 122–129
Monoculodes edwardsi, 50
Montagna, P. A., and E. Ruber, 61–64
Morone
  americana, 38, 139–140
  saxatilis, 133, 140
Mugil
  cephalus, 93–96, 282
  curema, 93–96, 100–104
  spp., 99–100
Murawski, S. A., G. R. Clayton, R. J. Reed, and C. F. Cole, 308–314
Murphy, D. J., and E. McCausland, 318–320
Mussel, ribbed, 84
Mya arenaria, 84, 87
Myrophis punctatus, 89, 93–96
Myriophyllum spicatum, fauna in beds of, 38–54
Mysidium columbiae, 321
Mysidopsis almyra, 321–323
  bahia, 321, 323
Mystic River Estuary, Conn., 1–10
Mytilopsis (=*Congeria*) *leucopheata*, 85–86

N

Nannochloris atomus, 213
Naphthalene, threshold for detection by blue crab, 224–229
Narragansett Bay, Rhode Island, 106–110
Navicula, 214, 275
Neanthes arenaceodentata, 228
Necropsy, of ospreys, 155–167
Neff, J. M., and L. A. Reitsema, 321–323
Nematodes, 296, 301
  density on decomposing *Spartina*, 61–64
Neoagardhiella baileyi, 200, 204
Neomysis, 321
  mercedis, 301
Neopanope sayi, 290
Nereis succinea, ingestion rate, 55–60
New York Bight, 75–83
Nitrogen, distribution and cycling in salt marsh, 111–121
Nitzschia
  incrustans, 125
  pungens, 208
  seriata, 213
  sigma, 271, 272, 274–276
  vermicularis, 125
North Inlet estuary, South Carolina, 90, 142–147, 236–241
Notemigonus chrysoleucas, 185
Notropis hudsonius, 39

O

Old Plantation Creek, Virginia, 207–216
Oligochaetes, 296, 303
Olla, B. L., and W. H. Pearson, 224–229
Oncorhynchus
  gorbuscha, 313
  kets, 313
  kisutch, 140
Organochlorines, in ospreys, 155–167
Orth, R. J., and K. L. Heck, Jr., 278–288, 289–295
Orris, P. K., 200–206
Osmerus mordax, 308–314
Osprey, pollutants in, 155–167
Oxygen, seasonal depletion in Chesapeake Bay, 242–247
Oyster, 146, 186, 189

P

Pachygrapsus, 127
  crassipes, 228
Pagurus longicarpus, 290
Palaemon macrodactylus, 301
Palaemonetes
  intermedius, 290, 293
  pugio, 133, 184, 186–188, 198, 228, 290, 292
  spp., 289
  vulgaris, 290
Panaeus aztecus, 228, 290–291, 293
Pandion haliaetus, 155–167
Panopeus spp., 220
Parahaustorius attenuatus, 79
Paralichthys
  albiguttata, 94–96
  dentatus, 93–96, 280
  lethostigma, 94–96
Paravortex
  cardii, 84
  gemellipara, 84–88
  scrobiculariae, 84
Parhyale hawaiensis, 228
Parker River, Massachusetts, 308–314
Patrick, W. H., Jr., *see* Buresh, R. J.
PCB, 155–156, 159
Pearson, W. H., and B. L. Olla, 224–229
Pelecanus
  occidentalis, 162
  onocrotalus, 164
Peloscolex benedeni, 49
Periclimenes, 293
Phasianus colchicus, 165
Phoeniconaias minor, 164

Phosphorus, distribution and cycling in salt marsh, 111–121
Photosynthesis, 315–317
Photosynthesis inhibitor, 217–223, 271–277
Phragmites communis, 174
Phytoplankton
  release of dissolved organic carbon, 230–233
  seasonal composition, 207–216
Pimephales promelas, 133, 142
Plantago maritima, 174
Pleuronectes platessa, 104
Plumley, F. G.,
  D. E. Davis, J. T. McEnerney, and J. W. Everest, 217–223
  and D. E. Davis, 271–277
Pollutants in ospreys, 155–167
Polychaete, nocturnal swimming, 148–149
Polypedilum
  digitifer, 44–45, 48–49
  illinoense, 44–45, 47–48
Polyploidy, in *Spartina alterniflora*, 168–170
Polysiphonia, 200, 203–205
  denudata, 204
  elongata, 204
  fibrillosa, 204
  flexicaulis, 204
  harveyi, 204
  nigrescens, 204
  subtilissima, 204
Pomatomus saltatrix, 282
Pondweed, 173
Porphyra sp., 200, 204
Potamogeton
  crispus, 52
  pectinatus, 52
  perfoliatus, 52
  spp., 173
Potomac River estuary, 11–19, 150–151
Primary productivity, 242–247
Prionotus tribulus, 92
Procladius
  bellus, 50
  sublettei, 44–45, 48–49
Productivity
  of algal epiphytes, 315–317
  of salt marsh algal mat, 122–131
Prorocentrum micans, 213
Protohaustorius
  deichmannae, 79, 83
  wigleyi, 79, 83

R

Rabenold, G. E., and D. G. Cargo, 20–27
Recruitment *Crangon septemspinosa*, 1–10
Reed, R. J., and R. B. Dormermuth, 65–68
Reed, R. J., *see* Murawski, S. A.
Reitsema, L. A., and J. M. Neff, 321–323
Reproduction, asexual, of *Chrysaora quinquecirrha*, 20–27
Respiration, in stratified water, 242–247
Rhabdocoele flatworm, 84–88
Rhabdonema minutum, 214
Rheotanytarsus sp., 44, 48
Rhepoxynius epistomus, 79
Rhithropanopeus harrisi, 42, 50
Rhizoclonium riparium, 125
Rhizosolenia,
  alata, 215
  calcar avis, 208, 215
  fragilissima, 215
  stolterfothii, 215
  styliformis, 215
Rhodomela confervoides, 204
Ritchie, D. E., book reviews, 152–153
Roberts, M. H., Jr., 184–191
  and A. T. Leggett, Jr., 192–199
Rotenone collection efficiency, 98–105
Ruber, E., and P. A. Montagna, 61–64
Rudlow, A., 177–183
Ruppia maritima, 278, 290

S

Sacramento River, California, 296–307
Sagitta elegans, 106–110
Sagittaria
  graminea, 174
  variabilis, 173
Saila, S. B., book reviews, 325
Salicornia bigelovii, 124
Salinity
  distribution and tidal mixing, 248–257
  response of benthos to changes in, 296–307
  variation on tidal flat, 28–37
Salmo gairdneri, 133, 140
Salt marsh cord grass, 61–64, 111–121, 168–176
Salt marsh productivity, 315–317
Salvelinus fontinalis, 142
San Francisco Bay Estuary, California, 296–307
Schizothrix, 125
Schwing, F. B., and B. Kjerfve, 236–241
Sciaenops ocellata, 282
Scolecolepides viridis, 42, 148–149
Scrobicularia plana, 84
Sea nettle, asexual reproduction, 20–27
Sediments
  deposition on tidal flat, 28–37
  ingestion rates by deposit feeders, 55–60
  labeling with fluorescent particles, 55–60
  response of benthos to changes in, 296–307
Seine collection efficiency, 98–105
Seirospora, 204
Sesarma cinereum, 218, 220
Shifrin, N. S., 230–233
Shrimp
  brown, 291
  grass, 133
  mysid, 321–323
  opossum, 321–323
  palaemonid, 289
  sand, 1–10, 289
Siegfried, C. A., M. E. Kopache, and A. W. Knight, 296–307
Skeletonema costatum, 207–208, 213–215
Spartina
  alterniflora, 90, 92, 130, 217–219, 236, 240, 266–269, 271, 315–317
    decomposition rate, 61–64
    height, 168–176
    polyploidy, 168–170
    utilization of nitrogen and phosphorus, 111–121
  angelica, 169
  cynosuroides, 240
  foliosa, 122–130
  maritima, 169
  × *neyrautii*, 169
  patens, 266
  pectinata, 169
  spartinae, 169
  stricta, 169
  × *townsendii*, 169
Spawning movements of *Osmerus mordax*, 308–314
Spirinchus thaleichthys, 309
Spongomorpha arcta, 205
Spyridia, 205
Stratification, of oxygen, 242–247
Stream channel migration, 258–270
Strix aluco, 164
Strongylura marina, 282
Suaeda depressa, 173
Sweatt, A. J., 106–110
Swimming, nocturnal of *Scolecolepides viridis*, 148–149
Symphurus plaguisa, 92
Syngnathus
  floridae, 280
  fuscus, 278, 280
  louisianae, 92

T

Tachypleus gigas, 181–182
Taft, J. L., W. R. Taylor, E. O. Hartwig, and R. Loftus, 242–247
Tagging studies, of *Osmerus mordax*, 308–314
Tanytarsus sp., 44–47, 49
Tautoga onitis, 280
Taylor, W. R., *see* Taft, J. L.
Temperature acclimation, 318–320
Temperature,
  effect on sediment settling rates on tidal flat, 28–37
Thalassionema nitzschioides, 212, 215
Thalassiosira fluviatilis, 271, 272, 274–276

Thalassiosira nordemskioldi, 208, 212, 215
Thomson, B. E., R. C. Worrest and H. Van Dyke, 69–72
Thor, 293
Tidal flat, sedimentation rate on, 28–37
Tidal mixing and salinity distribution, 248–257
Tidal stream channel migration and morphology, 258–270
Tijuana Estuary, California, 122–131
Tourtellotte, G. H., *see* Dauer, D. M.
Toxaphene, in rain, 142–147
Toxicity, of atrazine, 217–223
Tozeuma, 293
Trachyneis aspera, 125
Trapa natans, 52
Tubifex tubifex, 51
Typha latifolia, 173–174
Tyto alba, 164

U

Uca, 127
  pugilator, 221–222
  Pugnax, 217–223, 272–273
Ulva, 200, 205
Urospora sp., 200, 205
Ultraviolet radiation, 69–72

V

Vallisneria americana, 52
Van Dyke, H., *see* Thomson, B. E.
Vaucheria sp., 200, 204
Vegetation, influence on erosion in tidal marshes, 258–270

W

Wardle, W. J., 84–88
Waterhyacinth, 173
Waterplantain, 173
Water properties of tidal flat, 28–37
Wiemeyer, S. N., T. G. Lamont, and L. N. Locke, 155–167
Weinstein, M. P., and R. W. Davis, 98–105
Wigeon grass, 278, 290
Wilson, R. E., and A. D. Hamilton, 11–19
Winyah Bay, South Carolina, 236–241
Worrest, R. C., *see* Thomson, B. E.

Z

Zedler, J. B., 122–131
Zinc, 155, 157, 163
Zingmark, R., book review, 324–325
Zizania aquatica, 267
Zostera marina meadows
  decapod crustacea in, 289–295
  fishes in, 278–288